

Biological switches and clocks

John J Tyson, Reka Albert, Albert Goldbeter, Peter Ruoff and Jill Sible

J. R. Soc. Interface 2008 **5**, S1-S8
doi: 10.1098/rsif.2008.0179.focus

References

This article cites 102 articles, 35 of which can be accessed free

http://rsif.royalsocietypublishing.org/content/5/Suppl_1/S1.full.html#ref-list-1

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

To subscribe to *J. R. Soc. Interface* go to: <http://rsif.royalsocietypublishing.org/subscriptions>

Biological switches and clocks

John J. Tyson^{1,*}, Reka Albert², Albert Goldbeter³,
Peter Ruoff⁴ and Jill Sible¹

¹Department of Biological Sciences, Virginia Polytechnic Institute & State University,
Blacksburg, VA 24061, USA

²Department of Physics, Pennsylvania State University, University Park, PA 16802, USA

³Department of Chemical Physics, Free University of Brussels, 1050 Brussels, Belgium

⁴Faculty of Science & Technology, University of Stavanger, 4036 Stavanger, Norway

To introduce this special issue on biological switches and clocks, we review the historical development of mathematical models of bistability and oscillations in chemical reaction networks. In the 1960s and 1970s, these models were limited to well-studied biochemical examples, such as glycolytic oscillations and cyclic AMP signalling. After the molecular genetics revolution of the 1980s, the field of molecular cell biology was thrown wide open to mathematical modellers. We review recent advances in modelling the gene–protein interaction networks that control circadian rhythms, cell cycle progression, signal processing and the design of synthetic gene networks.

Keywords: bistability; oscillations; chemical reaction networks

1. INTRODUCTION

The living cell receives signals from its environment and its own internal state, processes the information, and initiates appropriate responses in terms of changes in gene expression, cell movement, and cell growth or death. Like a digital computer, information processing within cells is carried out by a complex network of switches and oscillators, but instead of being fabricated from silicon transistors and quartz crystals, the cell's computer is an evolved network of interacting genes and proteins. In the same way that computer design was made possible by a sophisticated theory of electronic circuitry, a basic understanding of cellular regulatory mechanisms will require a relevant theory of biomolecular circuitry. Although the 'engineering mindset' is sorely needed to make sense of the cell's circuitry, the squishy, sloppy, massively parallel, analogue nature of biochemistry is so different from the solid-state, precise, sequential, digital nature of computers that the mathematical tools and intellectual biases of the solid-state physicist/electrical engineer are not entirely appropriate to unravelling the molecular logic of cell physiology. New modelling paradigms and software tools are evolving to meet the challenges of the new 'systems biology' of the living cell.

In this context, we organized a six-week workshop on 'Biological switches and clocks' at the Kavli Institute for Theoretical Physics in Santa Barbara, CA, in summer 2007. The goal of the workshop was to bring together a diverse group of theorists and experimentalists, who

shared a common interest in understanding the molecular mechanisms that control the physiological properties of living organisms. The programme was organized around a few main themes: circadian rhythms; signalling networks; cell cycle regulation; synthetic gene networks; and deterministic and stochastic modelling. This special issue of *Interface* comprised peer-reviewed contributions from the participants in the workshop.

2. HISTORICAL CONTEXT

The first 'theoretical biologists' to consider biological interactions as dynamical systems were Alfred Lotka (a physical chemist) and Vito Volterra (a mathematician), who studied the pair of nonlinear ordinary differential equations

$$\frac{dx}{dt} = f(x, y), \quad \frac{dy}{dt} = g(x, y), \quad (2.1)$$

where x and y are the variables describing the time-dependent state of a chemical or biological system. Lotka (1924) and Volterra (1931) proposed models with periodic solutions (clocks) and multiple steady states (switches), which Volterra applied in the field of population biology to describe predator–prey oscillations and competition between species. For example, the Lotka–Volterra equations for prey (x) and predator (y) species are

$$\frac{dx}{dt} = ax - bxy, \quad \frac{dy}{dt} = cxy - dy, \quad (2.2)$$

where a , b , c , ... are 'rate' constants. Equations (2.2) have a one-parameter family of nested periodic solutions given by a 'conservation of energy' condition

$$E(x, y) = a \ln y - by + d \ln x - cx = \text{const.}$$

*Author for correspondence (tyson@vt.edu).

One contribution of 10 to a Theme Supplement 'Biological switches and clocks'.

The Lotka–Volterra equations for two competing species (x and y) are

$$\left. \begin{aligned} \frac{dx}{dt} &= ax(K-x) - bxy, \\ \frac{dy}{dt} &= cy(L-y) - dxy. \end{aligned} \right\} \quad (2.3)$$

If $c/d < K/L < b/a$, then these differential equations have two stable steady-state solutions

$$(x = K, y = 0) \quad \text{and} \quad (x = 0, y = L),$$

separated by an unstable steady state at

$$\left(x = c \frac{bL - aK}{bd - ac}, y = a \frac{dK - cL}{bd - ac} \right).$$

Dynamical systems (2.2) and (2.3) are paradigms of oscillations and bistability in a biological context.

In equations (2.2) and (2.3), x and y are interpreted as populations of interacting biological species in some common physical environment, but, in principle, they could just as well be interpreted as concentrations of reacting biochemical moieties in a cell. In the latter interpretation, the systems would model biochemical oscillations and bistability. But in the 1930s, no one was seriously thinking along these lines, because there were no known examples of biochemical clocks or switches. The field of cell biology was only beginning to find its roots in biochemistry and genetics.

To be sure, circadian rhythms and spontaneous oscillatory electrical impulses were recognized as interesting and important cellular phenomena, but the molecular bases of these activities were completely unknown. Embryonic development also inspired speculations about how biochemical reactions might spontaneously break symmetry and generate spatial patterns, but, aside from a singular paper by Turing (1952), no one had any idea on how ‘biochemistry’ might escape the boring fate of uniform chemical equilibrium. Classical biochemistry, with its linear pathways converting substrates A and B into end products Y and Z, seemed to offer little scope for exotic chemical dynamics.

This situation changed dramatically in the 1950s and the early 1960s, with the recognition of positive and negative feedback control of gene expression (Jacob & Monod 1961; Umbarger 1961) and with the discoveries of bistability in *lac* operon induction (Novick & Weiner 1957), oscillations in yeast glycolysis (Chance *et al.* 1964; Ghosh & Chance 1964) and periodic enzyme synthesis in bacteria (Masters & Donachie 1966). Biochemists began to appreciate the importance of switches and clocks in molecular cell biology. In 1965, Brian Goodwin presented a simple model of periodic enzyme synthesis based on the negative feedback of metabolic end products on gene transcription (Goodwin 1965), and in 1968 Joseph Higgins published a ground-breaking paper on the theory of biochemical oscillations in enzyme-catalysed reaction networks (Higgins 1967). The work by Field *et al.* (1972) and Field & Noyes (1974) on the oscillatory Belousov–Zhabotinsky reaction provided the

first detailed and predictive mechanistic description of a reaction system showing bistability as well as temporal, spatial and chaotic oscillations (Field & Burger 1985; Epstein & Pojman 1998).

A particularly simple example of a chemical reaction network exhibiting exotic dynamics (oscillations, bistability, pattern formation, travelling waves) was the ‘Brusselator’ of Prigogine & Lefever (1968), whose temporal dynamics are governed by

$$\left. \begin{aligned} \frac{dx}{dt} &= a - bx + cx^2y - dx, \\ \frac{dy}{dt} &= bx - cx^2y - ey. \end{aligned} \right\} \quad (2.4)$$

For the special case $e=0$ (as originally proposed by Prigogine & Lefever), this dynamical system has a unique steady state at $(x=a/d, y=bd/ac)$, which is unstable if $d^3 - bd^2 + ca^2 < 0$. When the steady state is unstable, it is surrounded by a unique, stable limit cycle oscillation (figure 1). For $e \neq 0$, it is possible for the Brusselator to exhibit bistability, as illustrated in figure 2. The regions of oscillations and bistability come together in a characteristic X-shaped two-parameter bifurcation diagram (figure 3). Oscillations and bistability in the Brusselator are based on the ternary autocatalytic reaction, $Y + 2X \rightarrow 3X$. Exploiting the exotic properties of this reaction, Sel’kov (1968) presented a simple, effective model of glycolytic oscillations, and Schlögl (1972) devised a simple, popular model of bistability.

The early theoretical work of Goodwin, Higgins, Prigogine and Lefever, Sel’kov and Schlögl (Goodwin 1965; Higgins 1967; Prigogine & Lefever 1968; Sel’kov 1968; Schlögl 1972) convinced biophysical chemists that oscillations and bistability were definitely to be expected in biochemical reaction systems with positive and negative feedbacks to destabilize steady-state solutions. Although a ternary autocatalytic reaction such as $Y + 2X \rightarrow 3X$ is unlikely in physical chemistry, high-order nonlinearities in substrate concentration can be achieved by multi-subunit enzymes of the type being discovered at that time to regulate biochemical pathways (Monod *et al.* 1965; Goldbeter & Lefever 1972).

Bistability and oscillations were also recognized, at this time, as expected features of genetic regulatory systems with positive and negative feedbacks. A simple and elegant theoretical treatment of these cases was published by Griffith in 1968. For a positive feedback system ($x = \text{mRNA}$, $y = \text{protein}$),

$$\frac{dx}{dt} = \frac{a + y^2}{K^2 + y^2} - bx, \quad \frac{dy}{dt} = cx - dy, \quad (2.5)$$

Griffith showed that mRNA level can easily show bistable (low or high) expression as a function of the demand d for the protein (Griffith 1968*a*). In an accompanying paper, Griffith (1968*b*) showed that the corresponding negative feedback system,

$$\frac{dx}{dt} = \frac{a}{K^p + y^p} - bx, \quad \frac{dy}{dt} = cx - dy, \quad (2.6)$$




Figure 1. Oscillations in the Brusselator equation (2.4). Parameter values: $a=1$, $b=3$, $c=1$, $d=1$ and $e=0$. (a) Phase plane. Dashed lines are the x -nullcline, $y=[(b+d)x-a]/(cx^2)$, and the y -nullcline, $y=b/(cx)$. Solid line is the limit cycle oscillation, which encircles the unstable steady state (open circle). (b) One-parameter bifurcation diagram. Thick solid line is a locus of stable steady-state solutions, interrupted by (thin solid line) a locus of unstable steady-state solutions. The circles indicate how the amplitude of the limit cycle oscillation (maximum and minimum values of x) depends on d between the two Hopf bifurcation (HB) points, at $d=0.6527$ and 2.879 .

has a unique, stable steady state (i.e. no bistability or oscillations). However, a negative feedback loop with n components,

$$\left. \begin{aligned} \frac{dx_1}{dt} &= \frac{a_0}{K^p + x_n^p} - b_1 x_1, \\ \frac{dx_2}{dt} &= a_1 x_1 - b_2 x_2, \\ &\dots \\ \frac{dx_n}{dt} &= a_{n-1} x_{n-1} - b_n x_n, \end{aligned} \right\} \quad (2.7)$$

can exhibit limit cycle oscillations, as Griffith proved, if the nonlinearity p of the feedback control is sufficiently strong: $p > \sec^n(\pi/n)$. These compelling results laid the foundation for a theory of genetic regulatory systems (Tyson & Othmer 1978).

3. INTERLUDE

Although these scientists in the 1960s and 1970s created a mathematical theory of switches and clocks




Figure 2. Bistability in the Brusselator equation (2.4). Parameter values: $a=1$, $b=3$, $c=1$, $d=0.2$ and $e=0.4$. (a) Phase plane. Dashed lines are the x - and y -nullclines as shown in figure 1. Solid circles, stable steady states; open circle, unstable steady state (saddle point). (b) One-parameter bifurcation diagram. The solid line is the locus of steady-state solutions. The upper and lower branches are stable steady states; the intermediate branch tracks unstable steady states. The stable and unstable branches come together at saddle-node (SN) bifurcation points, at $d=0.07894$ and 0.2273 .




Figure 3. Two-parameter bifurcation diagram for the Brusselator. The solid lines track the loci of SN bifurcations in dependence on d and e . Note that, for example, for $e=0.4$, two SNs occur at $d=0.07894$ and 0.2273 , as shown in figure 2b. The dashed lines track the HB points. For $e=0$, two HBs occur at $d=0.6527$ and 2.879 , as shown in figure 1b.

in molecular cell biology, there were only a few examples of interesting cell physiology for which enough molecular machinery was known to apply the

Table 1. Molecular cell biologists recognizing a need for mathematical modelling.

Hartwell <i>et al.</i> (1999)	‘The best test of our understanding of cells will be to make quantitative predictions about their behaviour and test them. This will require detailed simulations of the biochemical processes taking place within [cells]. ... We need to develop simplifying, higher-level models and find general principles that will allow us to grasp and manipulate the functions of [biochemical networks].’
Venter (1999)	‘If we hope to understand biology, instead of looking at one little protein at a time, which is not how biology works, we will need to understand the integration of thousands of proteins in a dynamically changing environment. A computer will be the biologist’s number one tool.’
Fraser & Harland (2000)	‘[R]esults to date show a dizzying array of signalling systems acting within and between cells. ... In such settings, intuition can be inadequate, often giving incomplete or incorrect predictions. ... In the face of such complexity, computational tools must be employed as a tool for understanding.’
Nurse (2000)	‘Perhaps a proper understanding of the complex regulatory networks making up cellular systems like the cell cycle will require a ... shift from common sense thinking. We might need to move into a strange more abstract world, more readily analyzable in terms of mathematics than our present imaginings of cells operating as a microcosm of our everyday world.’

theory profitably. Notable successes involved theories of glycolytic oscillations, cyclic AMP signaling, and calcium oscillations and waves. For details, consult the books by Goldbeter (1996) and Winfree (2001). But other important problems—such as the origins of circadian rhythms, the regulation of fruit fly development, the coordination of cell growth and division or the chemotactic response of motile bacteria—lay beyond the grasp of theorists, because nothing was known about the genes and proteins which controlled these processes. Theoretical developments floundered in the 1980s, awaiting that decade’s spectacular advances in molecular genetics. By the early 1990s, there were enough reliable clues about molecular mechanisms for mathematical cell biologists to polish up their equations and rev up their computers. Early successes included the works of Bray *et al.* (1993) on bacterial chemotaxis, Novak & Tyson (1993) on frog egg cell cycles, Goldbeter (1995) and Ruoff & Rensing (1996) on circadian rhythms in *Drosophila* and *Neurospora*, Reintz *et al.* (1995) on fruit fly morphogenesis and McAdams & Shapiro (1995) on bacterial circuits.

By 2000, a major shift in opinion was developing in favour of quantitative modelling in molecular cell biology. This shift was powered by a growing number of significant theoretical contributions (Arkin *et al.* 1998; Sharp & Reintz 1998; Van Dassow *et al.* 2000; Asthagiri & Lauffenburger 2001), an increasing recognition among molecular biologists of the field’s need for mathematical models (table 1) and the commitment of key administrators at NIH, NSF, DOE and elsewhere to target research support to quantitative biology.

4. MODERN DEVELOPMENTS

Since 2000, the field of molecular systems biology has blossomed, with the founding of an International Society for Systems Biology (www.iscb.org), of new institutes, centres and departments of systems biology across the world, and of high-profile journals (*Molecular Systems Biology*, *PLoS Computational Biology*). It is not practical here to review all the great works that have been published in recent years, but we will mention some developments that are particularly relevant to the KITP workshop on biological switches and clocks.

Perhaps the most compelling proof of the relevance of quantitative modelling in molecular cell biology was the

design of artificial genetic networks that function as toggle switches and oscillators, in Jim Collins’ laboratory at Boston University (Gardner *et al.* 2000) and Stan Leibler’s laboratory at Princeton University (Elowitz & Leibler 2000). Both circuits were designed first in computer models quite similar to equations (2.5) and (2.7) of Griffith. Building the circuits in bacteria was a standard exercise in genetic engineering; the tricky part was engineering the proteins to have the right kinetic rate constants for synthesis and degradation. The models predicted ranges for these rate constants that must be respected in order to observe the desired behaviour, and the circuits behaved exactly as predicted by the models. From these pioneering demonstrations has arisen a new and exciting technology called synthetic biology (Hasty *et al.* 2001).

An example of how modelling can be used to design gene regulatory networks of specific functionality is found in the article by Conrad *et al.* (2008) in this special issue. They are interested in networks that respond to an input pulse train as a ‘resonator’ (i.e. output signal only if the pulse train has a desired frequency) or as an ‘integrator’ (i.e. output signal only after a desired total number of pulses). For a variety of network topologies, they show that either functionality can be achieved, provided the kinetic rate constants in the model are chosen within an appropriate range of values. That is to say, functionality is tied not so much to topology as to rate constant values. Hence, in designing gene regulatory networks, it is essential to pick not only a suitable network topology but also to engineer the rate constants into the correct range for the desired behaviour.

Quantitative modelling has also had a major impact on our understanding of the molecular basis of circadian rhythms. As new genes and proteins have been discovered and their roles in circadian physiology worked out, mathematical models of the regulatory network have become more sophisticated and accurate (Leloup & Goldbeter 2000; Forger & Peskin 2003; Locke *et al.* 2005; Ruoff *et al.* 2005a). Because the physiology and molecular biology of circadian rhythms are sufficiently different in diverse organisms (bread moulds, plant cells, fruit flies, mammals), models must be tailored to specific organisms, while also laying bare some of the unifying principles of circadian control in eukaryotes. Circadian rhythms of cyanobacteria, on the other hand, are controlled by a completely different

molecular mechanism that has provided a challenging problem for mathematical modellers (Emberly & Wingreen 2006; Mehra *et al.* 2006; Mori *et al.* 2007; van Zon *et al.* 2007). Many crucial properties of circadian physiology, such as temperature compensation, synchronization and phase resetting, are inherently dynamical in nature and require precise theoretical descriptions to be understood and managed (Ruoff 1992; Rand *et al.* 2004; Stelling *et al.* 2004; Ruoff *et al.* 2005*b*; Kurosawa & Iwasa 2005; Bagheri *et al.* 2007; Hong *et al.* 2007). In this special issue, Bagheri *et al.* (2008) used systems-theoretic tools (mathematical control theory) to explore the phase response characteristics of a noisy circadian clock model, in order to understand how populations of oscillators entrain one another to generate a robust rhythm and how the rhythm re-synchronizes to an external 24 hours light–dark cycle from an initial 8 hours phase shift (say, flying between Chicago and Paris).

The molecular machinery controlling cell growth and division in yeast cells, embryos and mammalian somatic cells received a great deal of attention from molecular biologists in the 1990s, with a few theoretical groups trying to keep pace with the flood of genetic and biochemical data (Obeyesekere *et al.* 1996; Kohn 1998; Aguda & Tang 1999; Chen *et al.* 2000; Qu *et al.* 2004). Although the experimentalists paid little attention to modellers during this time, in recent years there have appeared many influential papers that self-consciously test (and confirm) predictions of the models (Cross *et al.* 2002, 2005; Cross 2003; Pomerening *et al.* 2003; Sha *et al.* 2003) or bring modelling to bear on experimental design and interpretation (Pomerening *et al.* 2005; Queralt *et al.* 2006). All eukaryotes use the same basic mechanism, based on cyclin-dependent kinases, to regulate the progression of DNA synthesis, mitosis and cell division (Csikasz-Nagy *et al.* 2006), but the idiosyncrasies of specific cell types require specifically tailored mathematical models (Chen *et al.* 2004; Calzone *et al.* 2007). The cell cycles of bacteria are regulated by a completely different set of genes and proteins, unrelated to the eukaryotic control system, and modellers have focused on specific details of the gene–protein interaction network in α -proteobacteria (Shen *et al.* 2007; Li *et al.* 2008), as well as convergent evolution of the dynamical properties of the two distinct control systems (Brazhnik & Tyson 2006).

Signalling networks have also provided a rich testing ground for mathematical modelling of molecular regulatory networks. The classic example, bistability in the *lac* operon, has received thorough attention over the years, as reviewed by Santillán & Mackey (2008) in this issue. In the late 1990s, Ferrell and colleagues published a series of papers on the MAP kinase pathway in *Xenopus* eggs, in which they experimentally demonstrated bistability of its response to progesterone signals, and used modelling to suggest that this bistability is due to positive feedback in the signalling pathway (Ferrell & Machelder 1998; Ferrell & Xiong 2001). Recently, Kholodenko and others have shown theoretically that positive feedback in the MAP kinase pathway is not necessary for bistability but can arise subtly from the multisite phosphorylation reactions that are ubiquitous

features of these kinase cascades (Markevich *et al.* 2004; Gunawardena 2005; Chickarmane *et al.* 2007). A major emphasis of the workshop was modelling of signalling networks, for example, pheromone signalling in yeast (Behar *et al.* 2007), bacterial chemotaxis (Keymer *et al.* 2006), regulatory circuits in the AIDS virus (Weinberger & Shenk 2007; Weinberger *et al.* 2008) and osmo-adaptation in yeast (Mettetal *et al.* 2008). In this special issue, the relevant contributions are by Csikasz-Nagy & Soyer (2008) on ‘adaptation’ in a simple biochemical network and by Jang & Gomer (2008) on size regulation in *Dictyostelium*.

In all these cases, the molecular reaction networks are known to be very complex and full of regulatory signals (feedback and feed-forward). The networks generate complex dynamical activity that is crucial to the survival, behaviour, development and reproduction of the living cell. Correlating the physiology of cells to the underlying molecular networks is no longer possible by intuitive biochemical reasoning, and mathematical models now play a central role in uncovering these correlations, framing our mechanistic hypotheses, and testing their implications. This new wave of realistic modelling and rigorous testing requires a new generation of computational tools for model building, model composition, simulation of real experimental protocols, comparison of simulations to a large array of experimental data, estimation of kinetic parameters from the data, statistically significant tests of model predictions, assessments of model sensitivity and robustness, and systematic analysis of mechanistic alternatives. Much progress has been made on this front, as demonstrated by the sophisticated tools now available to modellers.

- Virtual Cell, <http://www.nrcam.uchc.edu/>
- Copasi (Complex Pathway Simulator), <http://www.copasi.org/tiki-index.php>
- Systems Biology Workbench, <http://www.sys-bio.org/research/sbwintro.htm>
- E-Cell, <http://www.e-cell.org/ecell/>
- JigCell, <http://jigcell.biol.vt.edu/>
- Cell Designer, <http://www.celldesigner.org/>
- Silicon Cell, <http://homepages.cwi.nl/~gollum/SiC/>

Nonetheless, the computational challenges confronting the field are great, and much work is desperately needed in the area of tool development. In this regard, the paper by Rand (2008) in this special issue presents a sophisticated and promising new approach to the thorny problem of parameter estimation in complex network models with many parameters.

Modelling by discrete networks and stochastic processes were two other foci of the workshop. Discrete (Boolean) models, pioneered by Kauffman, Glass and Thomas (Kauffman 1969; Thomas 1973; Glass 1975), have been used to advantage recently in a variety of areas, including development (Albert & Othmer 2003; Chaves *et al.* 2005), immunology (Thakar *et al.* 2007) and cell cycle (Li *et al.* 2004; Faure *et al.* 2006; Davidich & Bornholdt 2008). This approach is nicely represented by two articles in this special issue. Chaves & Albert (2008) used Boolean models to study the effects of cell division on expression patterns of the segment-polarity

genes during *Drosophila* embryonic development, and Bornholdt (2008) reviewed the strengths and limitations of Boolean modelling of cell cycle regulation. Stochastic models of signalling networks (Ozbudak *et al.* 2002; Thattai & Van Oudenaarden 2004; Acar *et al.* 2008) and circadian rhythms (Forger & Peskin 2005; Yi & Jia 2005; Gonze & Goldbeter 2006) were discussed in some detail. In this special issue, Gonze *et al.* (2008) presented a stochastic version of their model of stress-induced oscillatory shuttling of Msn2 (a yeast transcription factor) between nucleus and cytoplasm.

Finally, the participants engaged in some lively discussions about if, when and how to introduce mathematical modelling into molecular cell biology classes at the undergraduate level. The majority opinion was that undergraduate students should be introduced to quantitative modelling and that this might reasonably be done with the aid of 'teaching modules' designed to complement standard textbooks. These ideas are elaborated in an article by Sible and Holmes ('Teaching at the interface: does computational cell biology fit in the undergraduate curriculum') to be published elsewhere.

5. LOOKING FORWARD

The KITP workshop brought together theoreticians and experimentalists to discuss successful collaborations, gripe about existing hurdles and forge new collaborations at the interface of math and cell biology. This special issue documents some of these success stories and problematic areas. The field of computational cell biology is here to stay, and the future looks very bright indeed (Goldbeter 2002).

REFERENCES

- Acar, M., Mettetal, J. T. & Van Oudenaarden, A. 2008 Stochastic switching as a survival strategy in fluctuating environments. *Nat. Genet.* **40**, 471–475. (doi:10.1038/ng.110)
- Aguda, B. D. & Tang, Y. 1999 The kinetic origins of the restriction point in the mammalian cell cycle. *Cell Prolif.* **32**, 321–335. (doi:10.1046/j.1365-2184.1999.3250321.x)
- Albert, R. & Othmer, H. G. 2003 The topology of the regulatory interactions predicts the expression pattern of the segment polarity genes in *Drosophila melanogaster*. *J. Theor. Biol.* **223**, 1–18. (doi:10.1016/S0022-5193(03)00035-3)
- Arkin, A., Ross, J. & McAdams, H. H. 1998 Stochastic kinetic analysis of developmental pathway bifurcation in phage lambda-infected *Escherichia coli* cells. *Genetics* **149**, 1633–1648.
- Asthagiri, A. R. & Lauffenburger, D. A. 2001 A computational study of feedback effects on signal dynamics in a mitogen-activated protein kinase (MAPK) pathway model. *Biotechnol. Prog.* **17**, 227–239. (doi:10.1021/bp010009k)
- Bagheri, N., Stelling, J. & Doyle III, F. J. 2007 Circadian phase entrainment via nonlinear model predictive control. *Int. J. Robust Nonlin. Control* **17**, 1555–1571. (doi:10.1002/rnc.1209)
- Bagheri, N., Taylor, S. R., Meeker, K., Petzold, L. R. & Doyle III, F. J. 2008 Synchrony and entrainment properties of robust circadian oscillators. *J. R. Soc. Interface* **5**, S17–S28. (doi:10.1098/rsif.2008.0045.focus)
- Behar, M., Hao, N., Dohlman, H. G. & Elston, T. C. 2007 Mathematical and computational analysis of adaptation via feedback inhibition in signal transduction pathways. *Biophys. J.* **93**, 806–821. (doi:10.1529/biophysj.107.107516)
- Bornholdt, S. 2008 Boolean network models of cellular regulation: prospects and limitations. *J. R. Soc. Interface* **5**, S85–S94. (doi:10.1098/rsif.2008.0132.focus)
- Bray, D., Bourret, R. B. & Simon, M. I. 1993 Computer simulation of the phosphorylation cascade controlling bacterial chemotaxis. *Mol. Biol. Cell* **4**, 469–482.
- Brazhnik, P. & Tyson, J. J. 2006 Cell cycle control in bacteria and yeast: a case of convergent evolution? *Cell Cycle* **5**, 522–529.
- Calzone, L., Thieffry, D., Tyson, J. J. & Novak, B. 2007 Dynamical modeling of syncytial mitotic cycles in *Drosophila* embryos. *Mol. Syst. Biol.* **3**, 131. (doi:10.1038/msb4100171)
- Chance, B., Hess, B. & Betz, A. 1964 DPNH oscillations in a cell-free extract of *S. carlsbergensis*. *Biochem. Biophys. Res. Commun.* **16**, 182–187. (doi:10.1016/0006-291X(64)90358-4)
- Chaves, M. & Albert, R. 2008 Studying the effect of cell division on expression patterns of the segment polarity genes. *J. R. Soc. Interface* **5**, S71–S84. (doi:10.1098/rsif.2007.1345.focus)
- Chaves, M., Albert, R. & Sontag, E. D. 2005 Robustness and fragility of Boolean models for genetic regulatory networks. *J. Theor. Biol.* **235**, 431–449. (doi:10.1016/j.jtbi.2005.01.023)
- Chen, K. C., Csikasz-Nagy, A., Gyorffy, B., Val, J., Novak, B. & Tyson, J. J. 2000 Kinetic analysis of a molecular model of the budding yeast cell cycle. *Mol. Biol. Cell* **11**, 369–391.
- Chen, K. C., Calzone, L., Csikasz-Nagy, A., Cross, F. R., Novak, B. & Tyson, J. J. 2004 Integrative analysis of cell cycle control in budding yeast. *Mol. Biol. Cell* **15**, 3841–3862. (doi:10.1091/mbc.E03-11-0794)
- Chickarmane, V., Kholodenko, B. N. & Sauro, H. M. 2007 Oscillatory dynamics arising from competitive inhibition and multisite phosphorylation. *J. Theor. Biol.* **244**, 68–76. (doi:10.1016/j.jtbi.2006.05.013)
- Conrad, E., Mayo, A. E., Ninfa, A. J. & Forger, D. B. 2008 Rate constants rather than biochemical mechanism determine behavior of genetic clocks. *J. R. Soc. Interface* **5**, S9–S15. (doi:10.1098/rsif.2008.0046.focus)
- Cross, F. R. 2003 Two redundant oscillatory mechanisms in the yeast cell cycle. *Dev. Cell* **4**, 741–752. (doi:10.1016/S1534-5807(03)00119-9)
- Cross, F. R., Archambault, V., Miller, M. & Klovstad, M. 2002 Testing a mathematical model of the yeast cell cycle. *Mol. Biol. Cell* **13**, 52–70. (doi:10.1091/mbc.01-05-0265)
- Cross, F. R., Schroeder, L., Kruse, M. & Chen, K. C. 2005 Quantitative characterization of a mitotic cyclin threshold regulating exit from mitosis. *Mol. Biol. Cell* **16**, 2129–2138. (doi:10.1091/mbc.E04-10-0897)
- Csikasz-Nagy, A. & Soyer, O. S. 2008 Adaptive dynamics with a single two-state protein. *J. R. Soc. Interface* **5**, S41–S47. (doi:10.1098/rsif.2008.0099.focus)
- Csikasz-Nagy, A., Battogtokh, D., Chen, K. C., Novak, B. & Tyson, J. J. 2006 Analysis of a generic model of eukaryotic cell-cycle regulation. *Biophys. J.* **90**, 4361–4379. (doi:10.1529/biophysj.106.081240)
- Davidich, M. I. & Bornholdt, S. 2008 Boolean network model predicts cell cycle sequence of fission yeast. *PLoS ONE* **3**, e1672. (doi:10.1371/journal.pone.0001672)
- Elowitz, M. B. & Leibler, S. 2000 A synthetic oscillatory network of transcriptional regulators. *Nature* **403**, 335–338. (doi:10.1038/35002125)
- Emberly, E. & Wingreen, N. S. 2006 Hourglass model for a protein-based circadian oscillator. *Phys. Rev. Lett.* **96**, 038 303. (doi:10.1103/PhysRevLett.96.038303)
- Epstein, I. R. & Pojman, J. A. 1998 *An introduction to nonlinear chemical dynamics*. New York, NY: Oxford University Press.

- Faure, A., Naldi, A., Chaouiya, C. & Thieffry, D. 2006 Dynamical analysis of a generic Boolean model for the control of the mammalian cell cycle. *Bioinformatics* **22**, e124–e131. (doi:10.1093/bioinformatics/btl210)
- Ferrell Jr, J. E. & Machelder, E. M. 1998 The biochemical basis of an all-or-none cell fate switch in *Xenopus* oocytes. *Science* **280**, 895–898. (doi:10.1126/science.280.5365.895)
- Ferrell Jr, J. E. & Xiong, W. 2001 Bistability in cell signaling: how to make continuous processes discontinuous, and reversible processes irreversible. *Chaos* **11**, 227–236. (doi:10.1063/1.1349894)
- Field, R. J. & Burger, M. 1985 *Oscillations and traveling waves in chemical systems*. New York, NY: John Wiley.
- Field, R. J. & Noyes, R. M. 1974 Oscillations in chemical systems: IV. Limit cycle behavior in a model of a real chemical reaction. *J. Chem. Phys.* **60**, 1877–1884. (doi:10.1063/1.1681288)
- Field, R. J., Korös, E. & Noyes, R. M. 1972 Oscillations in chemical systems: II. Thorough analysis of temporal oscillation in the bromate–cerium–malonic acid system. *J. Am. Chem. Soc.* **94**, 8649–8664. (doi:10.1021/ja00780a001)
- Forger, D. B. & Peskin, C. S. 2003 A detailed predictive model of the mammalian circadian clock. *Proc. Natl Acad. Sci. USA* **100**, 14 806–14 811. (doi:10.1073/pnas.2036281100)
- Forger, D. B. & Peskin, C. S. 2005 Stochastic simulation of the mammalian circadian clock. *Proc. Natl Acad. Sci. USA* **102**, 321–324. (doi:10.1073/pnas.0408465102)
- Fraser, S. E. & Harland, R. M. 2000 The molecular metamorphosis of experimental embryology. *Cell* **100**, 41–55. (doi:10.1016/S0092-8674(00)81682-7)
- Gardner, T. S., Cantor, C. R. & Collins, J. J. 2000 Construction of a genetic toggle switch in *Escherichia coli*. *Nature* **403**, 339–342. (doi:10.1038/35002131)
- Ghosh, A. & Chance, B. 1964 Oscillations of glycolytic intermediates in yeast cells. *Biochem. Biophys. Res. Commun.* **16**, 174–181. (doi:10.1016/0006-291X(64)90357-2)
- Glass, L. 1975 Classification of biological networks by their qualitative dynamics. *J. Theor. Biol.* **54**, 85–107. (doi:10.1016/S0022-5193(75)80056-7)
- Goldbeter, A. 1995 A model for circadian oscillations in the *Drosophila* period protein (PER). *Proc. R. Soc. B* **261**, 319–324. (doi:10.1098/rspb.1995.0153)
- Goldbeter, A. 1996 *Biochemical oscillations and cellular rhythms*. Cambridge, UK: Cambridge University Press.
- Goldbeter, A. 2002 Computational approaches to cellular rhythms. *Nature* **420**, 238–245. (doi:10.1038/nature01259)
- Goldbeter, A. & Lefever, R. 1972 Dissipative structures for an allosteric model: application to glycolytic oscillations. *Biophys. J.* **12**, 1302–1315.
- Gonze, D. & Goldbeter, A. 2006 Circadian rhythms and molecular noise. *Chaos* **16**, 026 110. (doi:10.1063/1.2211767)
- Gonze, D., Jacquet, M. & Goldbeter, A. 2008 Stochastic modelling of nucleocytoplasmic oscillations of the transcription factor Msn2 in yeast. *J. R. Soc. Interface* **5**, S95–S109. (doi:10.1098/rsif.2008.0141.focus)
- Goodwin, B. C. 1965 Oscillatory behavior in enzymatic control processes. *Adv. Enzyme Regul.* **3**, 425–438. (doi:10.1016/0065-2571(65)90067-1)
- Griffith, J. S. 1968a Mathematics of cellular control processes: II. Positive feedback to one gene. *J. Theor. Biol.* **20**, 209–216. (doi:10.1016/0022-5193(68)90190-2)
- Griffith, J. S. 1968b Mathematics of cellular control processes: I. Negative feedback to one gene. *J. Theor. Biol.* **20**, 202–208. (doi:10.1016/0022-5193(68)90189-6)
- Gunawardena, J. 2005 Multisite phosphorylation makes a good threshold but can be a poor switch. *Proc. Natl Acad. Sci. USA* **102**, 14 617–14 622. (doi:10.1073/pnas.0507322102)
- Hartwell, L. H., Hopfield, J. J., Leibler, S. & Murray, A. W. 1999 From molecular to modular cell biology. *Nature* **402**, C47–C52. (doi:10.1038/35011540)
- Hasty, J., McMillen, D., Isaacs, F. & Collins, J. J. 2001 Computational studies of gene regulatory networks: in numero molecular biology. *Nat. Rev. Genet.* **2**, 268–279. (doi:10.1038/35066056)
- Higgins, J. 1967 The theory of oscillating reactions. *Ind. Eng. Chem.* **59**, 18–62. (doi:10.1021/ie50689a006)
- Hong, C. I., Conrad, E. D. & Tyson, J. J. 2007 A proposal for robust temperature compensation of circadian rhythms. *Proc. Natl Acad. Sci. USA* **104**, 1195–1200. (doi:10.1073/pnas.0601378104)
- Jacob, F. & Monod, J. 1961 Genetic regulatory mechanisms in the synthesis of proteins. *J. Mol. Biol.* **3**, 318–356.
- Jang, W. & Gomer, R. H. 2008 Combining experiments and modelling to understand size regulation in *Dictyostelium discoideum*. *J. R. Soc. Interface* **5**, S49–S58. (doi:10.1098/rsif.2008.0067.focus)
- Kauffman, S. A. 1969 Metabolic stability and epigenesis in randomly constructed genetic nets. *J. Theor. Biol.* **22**, 437–467. (doi:10.1016/0022-5193(69)90015-0)
- Keymer, J. E., Endres, R. G., Skoge, M., Meir, Y. & Wingreen, N. S. 2006 Chemosensing in *Escherichia coli*: two regimes of two-state receptors. *Proc. Natl Acad. Sci. USA* **103**, 1786–1791. (doi:10.1073/pnas.0507438103)
- Kohn, K. W. 1998 Functional capabilities of molecular network components controlling the mammalian G1/S cell cycle phase transition. *Oncogene* **16**, 1065–1075. (doi:10.1038/sj.onc.1201608)
- Kurosawa, G. & Iwasa, Y. 2005 Temperature compensation in circadian clock models. *J. Theor. Biol.* **233**, 453–468. (doi:10.1016/j.jtbi.2004.10.012)
- Leloup, J.-C. & Goldbeter, A. 2000 Toward a detailed computational model for the mammalian circadian clock. *Proc. Natl Acad. Sci. USA* **100**, 7051–7056. (doi:10.1073/pnas.1132112100)
- Li, R., Long, T., Lu, Y., Ouyang, Q. & Tang, C. 2004 The yeast cell-cycle network is robustly designed. *Proc. Natl Acad. Sci. USA* **101**, 4781–4786. (doi:10.1073/pnas.0305937101)
- Li, S., Brazhnik, P., Sobral, B. & Tyson, J. J. 2008 A quantitative study of the division cycle of *Caulobacter crescentus* stalked cells. *PLoS Comput. Biol.* **4**, e9. (doi:10.1371/journal.pcbi.0040009)
- Locke, J. C., Southern, M. M., Kozma-Bognar, L., Hibberd, V., Brown, P. E., Turner, M. S. & Millar, A. J. 2005 Extension of a genetic network model by iterative experimentation and mathematical analysis. *Mol. Syst. Biol.* **1**, 55–63. (doi:10.1038/msb4100018)
- Lotka, A. J. 1924 *Elements of physical biology*. New York, NY: Dover. (Reprinted by Dover 1956.)
- Markevich, N. I., Hoek, J. B. & Kholodenko, B. N. 2004 Signaling switches and bistability arising from multisite phosphorylation in protein kinase cascades. *J. Cell Biol.* **274**, 36 362.
- Masters, M. & Donachie, W. D. 1966 Repression and control of cyclic enzyme synthesis in *Bacillus subtilis*. *Nature* **209**, 476–479. (doi:10.1038/209476a0)
- McAdams, H. H. & Shapiro, L. 1995 Circuit simulation of genetic networks. *Science* **269**, 650–656. (doi:10.1126/science.7624793)
- Mehra, A., Hong, C. I., Shi, M., Loros, J. J., Dunlap, J. C. & Ruoff, P. 2006 Circadian rhythmicity by autocatalysis. *PLoS Comput. Biol.* **2**, e96. (doi:10.1371/journal.pcbi.0020096)
- Mettetal, J. T., Muzzey, D., Gomez-Urbe, C. & van Oudenaarden, A. 2008 The frequency dependence of osmo-adaptation in *Saccharomyces cerevisiae*. *Science* **319**, 482–484. (doi:10.1126/science.1151582)

- Monod, J., Wyman, J. & Changeux, J. P. 1965 On the nature of allosteric transitions: a plausible model. *J. Mol. Biol.* **12**, 88–118.
- Mori, T., Williams, D. R., Byrne, M. O., Qin, X., Egli, M., Mchaourab, H. S., Stewart, P. L. & Johnson, C. H. 2007 Elucidating the ticking of an *in vitro* circadian clockwork. *PLoS Biol.* **5**, e93. (doi:10.1371/journal.pbio.0050093)
- Novak, B. & Tyson, J. J. 1993 Numerical analysis of a comprehensive model of M-phase control in *Xenopus* oocyte extracts and intact embryos. *J. Cell Sci.* **106**, 1153–1168.
- Novick, A. & Weiner, M. 1957 Enzyme induction as an all-or-none phenomenon. *Proc. Natl Acad. Sci. USA* **43**, 553–566. (doi:10.1073/pnas.43.7.553)
- Nurse, P. 2000 A long twentieth century of the cell cycle and beyond. *Cell* **100**, 71–78. (doi:10.1016/S0092-8674(00)81684-0)
- Obeyesekere, M. N., Herbert, J. R. & Zimmerman, S. O. 1996 A model of the G1 phase of the cell cycle incorporating cyclinE/cdk2 complex and retinoblastoma protein. *Oncogene* **11**, 1199–1205.
- Ozbudak, E. M., Thattai, M., Kurtser, I., Grossman, A. D. & Van Oudenaarden, A. 2002 Regulation of noise in the expression of a single gene. *Nat. Genet.* **31**, 69–73. (doi:10.1038/ng869)
- Pomerening, J. R., Sontag, E. D. & Ferrell Jr, J. E. 2003 Building a cell cycle oscillator: hysteresis and bistability in the activation of Cdc2. *Nat. Cell Biol.* **5**, 346–351. (doi:10.1038/ncb954)
- Pomerening, J. R., Kim, S. Y. & Ferrell Jr, J. E. 2005 Systems-level dissection of the cell-cycle oscillator: bypassing positive feedback produces damped oscillations. *Cell* **122**, 565–578. (doi:10.1016/j.cell.2005.06.016)
- Prigogine, I. & Lefever, R. 1968 Symmetry breaking instabilities in dissipative systems. II. *J. Chem. Phys.* **48**, 1695. (doi:10.1063/1.1668896)
- Qu, Z., Weiss, J. N. & MacLellan, W. R. 2004 Coordination of cell growth and cell division: a mathematical modeling study. *J. Cell Sci.* **117**, 4199–4207. (doi:10.1242/jcs.01294)
- Queralt, E., Lehane, C., Novak, B. & Uhlmann, F. 2006 Downregulation of PP2A(Cdc55) phosphatase by separase initiates mitotic exit in budding yeast. *Cell* **125**, 719–732. (doi:10.1016/j.cell.2006.03.038)
- Rand, D. A. 2008 Mapping global sensitivity of cellular network dynamics: sensitivity heat maps and a global summation law. *J. R. Soc. Interface* **5**, S59–S69. (doi:10.1098/2008.0084.focus)
- Rand, D. A., Shulgin, B. V., Salazar, D. & Millar, A. J. 2004 Design principles underlying circadian clocks. *J. R. Soc. Interface* **1**, 119–130. (doi:10.1098/rsif.2004.0014)
- Reintz, J., Mjolsness, E. & Sharp, D. H. 1995 Model for cooperative control of positional information in *Drosophila* by bicoid and maternal hunchback. *J. Exp. Zool.* **271**, 47–56. (doi:10.1002/jez.1402710106)
- Ruoff, P. 1992 Introducing temperature-compensation in any reaction kinetic oscillator model. *J. Interdiscipl. Cycle Res.* **23**, 92.
- Ruoff, P. & Rensing, L. 1996 The temperature-compensated Goodwin model simulates many circadian clock properties. *J. Theoret. Biol.* **179**, 275–285. (doi:10.1006/jtbi.1996.0067)
- Ruoff, P., Christensen, M. K. & Sharma, V. K. 2005a PER/TIM-mediated amplification, gene dosage effects and temperature compensation in an interlocking-feedback loop model of the *Drosophila* circadian clock. *J. Theor. Biol.* **237**, 41–57. (doi:10.1016/j.jtbi.2005.03.030)
- Ruoff, P., Loros, J. J. & Dunlap, J. C. 2005b The relationship between FRQ-protein stability and temperature compensation in the *Neurospora* circadian clock. *Proc. Natl Acad. Sci. USA* **102**, 17 681–17 686. (doi:10.1073/pnas.0505137102)
- Santillán, M. & Mackey, M. C. 2008 Quantitative approaches to the study of bistability in the *lac* operon of *Escherichia coli*. *J. R. Soc. Interface* **5**, S29–S39. (doi:10.1098/rsif.2008.0086.focus)
- Schlögl, F. 1972 Chemical reaction models for non-equilibrium phase transitions. *Z. Phys. A* **253**, 147–161.
- Sel'kov, E. E. 1968 Self-oscillations in glycolysis. 1. A simple kinetic model. *Eur. J. Biochem.* **4**, 79–86. (doi:10.1111/j.1432-1033.1968.tb00175.x)
- Sha, W., Moore, J., Chen, K., Lassaletta, A. D., Yi, C. S., Tyson, J. J. & Sible, J. C. 2003 Hysteresis drives cell-cycle transitions in *Xenopus laevis* egg extracts. *Proc. Natl Acad. Sci. USA* **100**, 975–980. (doi:10.1073/pnas.0235349100)
- Sharp, D. H. & Reintz, J. 1998 Prediction of mutant expression patterns using gene circuits. *Biosystems* **47**, 79–90. (doi:10.1016/S0303-2647(98)00014-8)
- Shen, X., Colleur, J., Shapiro, L., Dill, D., McAdams, H. & Horowitz, M. 2007 Hybrid modeling and robustness analysis on *Caulobacter* cell cycle regulation. In *Int. Conf. Systems Biology, 1–5 October 2007*. See www.stanford.edu/~xilings/Research/icsb2007.pdf.
- Stelling, J., Gilles, E. D. & Doyle III, F. J. 2004 Robustness properties of circadian clock architectures. *Proc. Natl Acad. Sci. USA* **101**, 13 210–13 215. (doi:10.1073/pnas.0401463101)
- Thakar, J., Pilione, M., Kirimanjeswara, B., Harvill, E. T. & Albert, R. 2007 Modeling systems-level regulation of host immune responses. *PLoS Comput. Biol.* **3**, e109. (doi:10.1371/journal.pcbi.0030109)
- Thattai, M. & Van Oudenaarden, A. 2004 Stochastic gene expression in fluctuating environments. *Genetics* **167**, 523–530. (doi:10.1534/genetics.167.1.523)
- Thomas, R. 1973 Boolean formalization of genetic control circuits. *J. Theor. Biol.* **42**, 563–585. (doi:10.1016/0022-5193(73)90247-6)
- Turing, A. M. 1952 The chemical basis of morphogenesis. *Phil. Trans. R. Soc. B* **237**, 37–72. (doi:10.1098/rstb.1952.0012)
- Tyson, J. J. & Othmer, H. G. 1978 The dynamics of feedback control circuits in biochemical pathways. *Prog. Theor. Biol.* **5**, 1–62.
- Umbarger, H. E. 1961 Feedback control by endproduct inhibition. *Cold Spring Harb. Symp. Quant. Biol.* **26**, 301–312.
- Van Dassow, G., Meir, E., Mumro, E. M. & Odell, G. M. 2000 The segment polarity network is a robust developmental module. *Nature* **406**, 188–192. (doi:10.1038/35018085)
- van Zon, J. S., Lubensky, D. K., Altena, P. R. H. & ten Wolde, P. R. 2007 An allosteric model of circadian KaiC phosphorylation. *Proc. Natl Acad. Sci. USA* **104**, 7420–7425. (doi:10.1073/pnas.0608665104)
- Venter, C. 1999 Computing 2010: from black holes to biology. *Nature* **402**, C67–C70. (doi:10.1038/35011561)
- Volterra, V. 1931 *Lecons sur la theorie de la lutte pour la vie*. Paris, France: Gauthier-Villars.
- Weinberger, L. S. & Shenk, T. 2007 An HIV feedback resistor: auto-regulatory circuit deactivator and noise buffer. *PLoS Biol.* **5**, e9. (doi:10.1371/journal.pbio.0050009)
- Weinberger, L. S., Dar, R. D. & Simpson, M. L. 2008 Transient-mediated fate determination in a transcriptional circuit of HIV. *Nat. Genet.* **40**, 466–470. (doi:10.1038/ng.116)
- Winfree, A. T. 2001 *The geometry of biological time*, 2nd edn. New York, NY: Springer.
- Yi, M. & Jia, Y. 2005 Light-noise-induced suprathreshold circadian oscillations and coherent resonance in *Drosophila*. *Phys. Rev. E* **72**, 012 902. (doi:10.1103/PhysRevE.72.012902)